ELSEVIER

Carbohydrate Research 283 (1996) C7–C8

CARBOHYDRATE RESEARCH

Contents

Regular papers

AM1 calculations on inclusion complexes of cyclomaltoheptaose (β-cyclodextrin) with 1,7-dioxaspiro[5.5]undecane and nonanal, and comparison with experimental results
A. Botsi, K. Yannakopoulou, E. Hadjoudis and J. Waite (Athens, Greece) 1

Tin(IV)-functionalised polymer supports; non-toxic and practical reagents for regioselective acetylation of sucrose
W.M. Macindoe (Tokyo, Japan), A. Williams (Canterbury, UK) and R. Khan (Trieste, Italy) 17

Syntheses of 1-*O*-carboxyalkyl GLA-60 analogues
M. Shiozaki, N. Deguchi, W.M. Macindoe, M. Arai, H. Miyazaki, T. Mochizuki, T. Tatsuta, J. Ogawa, H. Maeda and S.-i. Kurakata (Tokyo, Japan) 27

Characterization and interconversions of 2-*S*-ethyl-2-thio-D-mannose diethyl dithioacetal and the facile epimerization of 2-thio-D-mannopyranose derivatives
D. Horton, P. Norris and B. Berrang (Washington, DC, USA) 53

A nitro sugar derivative route to 2-thioepisophorose and 2-thiosophorose and their remarkable facile epimerization
M. Petrušová, E. Lattová, M. Matulová, L. Petruš (Bratislava, Slovak Republic) and J.N. BeMiller (West Lafayette, IN, USA) 73

Syntheses of triglycosyl tetrapeptides and a hexaglycosyl tetrapeptide
T. Takeda (Tokyo, Japan), T. Kanemitsu (Nagoya, Japan), N. Shimizu (Tokyo, Japan), Y. Ogihara (Nagoya, Japan) and M. Matsubara (Kobe, Japan) 81

Structure of the *O*-deacetylated glucuronoxylomannan from *Cryptococcus neoformans* Cap70 as determined by 2D NMR spectroscopy
B.E. Bacon, R. Cherniak (Atlanta, GA, USA), K.J. Kwon-Chung (Bethesda, MD, USA) and E.S. Jacobson (Richmond, VA, USA) 95

Structural studies of the lipopolysaccharide O-antigen and capsular polysaccharide of *Vibrio anguillarum* serotype O:2
I. Sadovskaya, J.-R. Brisson, E. Altman (Ottawa, Canada) and L.M. Mutharia (Guelph, Canada) . . 111

Structure of a decasaccharide isolated by mild acid degradation and dephosphorylation of the lipopolysaccharide of *Pseudomonas fluorescens* strain ATCC 49271
Y.A. Knirel (Borstel, Germany), J.H. Helbig (Dresden, Germany) and U. Zähringer (Borstel, Germany) 129

Acid modification of starch granules in alcohols: reactions in mixtures of two alcohols combined in different ratios
J.F. Robyt, J.-y. Choe, J.D. Fox, R.S. Hahn and E.B. Fuchs (Ames, IA, USA) 141

A periodic distribution of the chain length of amylopectin as revealed by high-performance anion-exchange chromatography
I. Hanashiro, J.-i. Abe and S. Hizukuri (Kagoshima, Japan) 151

Cell-wall polysaccharides from the marine green alga *Ulva "rigida"* (Ulvales, Chlorophyta) — NMR analysis of ulvan oligosaccharides
M. Lahaye and B. Ray (Nantes, France). 161

Structural study on a lipopolysaccharide from *Coxiella burnetii* strain Nine Mile in avirulent phase II
R. Toman and L. Škultéty (Bratislava, Slovak Republic) 175

Failure of glucose-binding lectins Con A and Lentil Lectin to identify glycation of haemoglobin
F. Frantzen (Mørkved, Norway) and E. Sundrehagen (Oslo, Norway) 187

Notes

NMR analysis of galactoglucan from *Pseudomonas marginalis*: assignment of the ^{1}H and ^{13}C NMR spectra and location of succinate groups
M. Matulová, L. Navarini (Trieste, Italy), S.F. Osman and W.F. Fett (Philadelphia, PA, USA). . . . 195

Polymerization of 1,2-anhydro-3,4-di-*O*-benzyl-β-L-rhamnopyranose
Q. Chen and F. Kong (Beijing, People's Republic of China) 207

Structural investigation of a cell-wall galactomannan from *Neurospora crassa* and *N. sitophila*
J.A. Leal, J. Jiménez-Barbero, B. Gómez-Miranda, A. Prieto, J. Domenech and M. Bernabé (Madrid, Spain). 215

NMR analysis of the structure of the O88 polysaccharide (O88 antigen) of *Escherichia coli* O88:K^{-}:H25
V.I. Torgov, A.S. Shashkov, H. Kochanowski, B. Jann and K. Jann (Freiburg, Germany) 223

Author index C1

Subject index C3

Contents C7